

## THEMED ISSUE: CANNABINOIDS

## RESEARCH PAPER

JWH018, a common constituent of 'Spice' herbal blends, is a potent and efficacious cannabinoid CB<sub>1</sub> receptor agonistBrady K Atwood<sup>1,2</sup>, John Huffman<sup>3</sup>, Alex Straiker<sup>1</sup> and Ken Mackie<sup>1</sup>

<sup>1</sup>The Gill Center and the Department of Psychological & Brain Sciences, Indiana University, Bloomington, IN, USA, <sup>2</sup>University of Washington, Program of Neurobiology and Behavior, Seattle, WA, USA, and <sup>3</sup>The Department of Chemistry, Clemson University, Clemson, SC, USA

**Background and purpose:** 'Spice' is an herbal blend primarily marketed in Europe as a mild hallucinogen with prominent cannabis-like effects and as a legal alternative to cannabis. However, a recent report identified a number of synthetic additives in samples of 'Spice'. One of these, the indole derivative JWH018, is a ligand for the cannabinoid receptor 1 (CB<sub>1</sub>) cannabinoid receptor and inhibits cAMP production in CB<sub>1</sub> receptor-expressing CHO cells. Other effects of JWH018 on CB<sub>1</sub> receptor-mediated signalling are not known, particularly in neurons. Here we have evaluated the signalling pathways activated by JWH018 at CB<sub>1</sub> receptors.

**Experimental approach:** We investigated the effects of JWH018 on neurotransmission in cultured autaptic hippocampal neurons. We further analysed its activation of ERK1/2 mitogen activated protein kinase (MAPK) and internalization of CB<sub>1</sub> receptors in HEK293 cells stably expressing this receptor.

**Key results:** In cultured autaptic hippocampal neurons, JWH018 potently inhibited excitatory postsynaptic currents ( $IC_{50} = 14.9$  nM) in a concentration- and CB<sub>1</sub> receptor-dependent manner. Furthermore, it increased ERK1/2 MAPK phosphorylation ( $EC_{50} = 4.4$  nM). We also found that JWH018 potently induced rapid and robust CB<sub>1</sub> receptor internalization ( $EC_{50} = 2.8$  nM;  $t_{1/2} = 17.3$  min).

**Conclusions and implications:** JWH018, a prominent component of several herbal preparations marketed for their psychoactivity, is a potent and effective CB<sub>1</sub> receptor agonist that activates multiple CB<sub>1</sub> receptor signalling pathways. Thus, it is likely that the subjective effects of 'Spice' are due to activation of cannabinoid CB<sub>1</sub> receptors by JWH018, added to this herbal preparation.

*British Journal of Pharmacology* (2010) **160**, 585–593; doi:10.1111/j.1476-5381.2009.00582.x; published online 22 January 2010

This article is part of a themed issue on Cannabinoids. To view the editorial for this themed issue visit <http://dx.doi.org/10.1111/j.1476-5381.2010.00831.x>

**Keywords:** Spice; JWH018; cannabinoid; CB<sub>1</sub> receptor; internalization; ERK1/2; MAPK; neurotransmission; marijuana; THC

**Abbreviations:** CB<sub>1</sub> receptor, cannabinoid receptor 1; CP47,497, 2-[(1R,3S)-3-hydroxycyclohexyl]-5-(2-methyloctan-2-yl)phenol; EPSC, excitatory postsynaptic current; ERK, extracellular signal regulated kinase; GPCR, G protein coupled receptor; JWH018, naphthalen-1-yl-(1-pentylindol-3-yl)methanone; MAPK, mitogen activated protein kinase; rimonabant, SR141716/5-(4-chlorophenyl)-1-(2,4-dichloro-phenyl)-4-methyl-N-(piperidin-1-yl)-1H-pyrazole-3-carboxamide; THC,  $\Delta^9$ -tetrahydrocannabinol; WIN55212-2, (R)-(+)-[2,3-dihydro-5-methyl-3-(4-morpholinylmethyl)pyrrolo[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenylmethanone mesylate

## Introduction

*Cannabis sativa* (cannabis, marijuana or hashish) is a widely used drug with well-known psychoactivity as well as potential medicinal value.  $\Delta^9$ -tetrahydrocannabinol (THC) has been identified as the principal psychoactive component of

Correspondence: Ken Mackie, Department of Psychological and Brain Sciences, Indiana University, 1101 E. 10th Street, Bloomington, IN 47405, USA. E-mail: [kmackie@indiana.edu](mailto:kmackie@indiana.edu)

Received 22 July 2009; revised 9 September 2009; accepted 5 October 2009

*C. sativa*, although it is only one of a number of bioactive phytocannabinoids found in the plant (Taura *et al.*, 2007). The physiological effects of THC have been well described (Ameri *et al.*, 1999; Howlett, 2002; Howlett *et al.*, 2002; Costa, 2007; Pertwee, 2008).

The cannabinoid receptor 1 (CB<sub>1</sub>) cannabinoid receptor (nomenclature follows Alexander *et al.*, 2008) has been identified as the receptor that mediates the behavioural effects of THC in animals (Monory *et al.*, 2007) and likely does so in humans (Huestis *et al.*, 2001). The CB<sub>1</sub> receptor is predominantly expressed in the CNS, particularly in areas such as the hippocampus, basal ganglia, cortex, amygdala and cerebellum – areas linked to behaviours affected by THC (Mackie, 2005). At the subcellular level, CB<sub>1</sub> receptors are primarily found on axon terminals, a prime location to influence neurotransmission (Gulyas *et al.*, 2004; Mackie, 2005; Nyiri *et al.*, 2005; Yoshida *et al.*, 2006; Matyas *et al.*, 2007). The CB<sub>1</sub> receptor is a G protein coupled receptor (GPCR) that couples to the G<sub>i/o</sub> class of G proteins and as such, upon agonist activation leads to an inhibition of adenylyl cyclase and subsequent decrease in cellular cAMP levels. CB<sub>1</sub> receptor activation also inhibits voltage gated calcium channels and activates inwardly rectifying potassium channels (Mackie *et al.*, 1995; Twitchell *et al.*, 1997; Howlett *et al.*, 2002). Cumulatively, these effects on intracellular signalling result in reduced cellular excitability and, due to its proximity to synaptic terminals (Nyiri *et al.*, 2005), in a reduction in the probability of neurotransmitter release (Shen *et al.*, 1996). This ability to inhibit neurotransmission allows both exogenous cannabinoid agonists (such as THC) and endogenous cannabinoids (endocannabinoids) to have a profound impact on neuronal communication. As CB<sub>1</sub> receptors are found on both glutamatergic and GABAergic terminals, their activation can suppress both inhibitory and excitatory synaptic transmission (Kreitzer and Regehr, 2001; Ohno-Shosaku *et al.*, 2001; Wilson and Nicoll, 2001; Chevaleyre *et al.*, 2006; Straiker and Mackie, 2006).

CB<sub>1</sub> receptor stimulation also results in activation of mitogen activated protein kinases (MAPKs), particularly extracellular signal-regulated kinase (p42/44 or ERK1/2) (Bouaboula *et al.*, 1995; Daigle *et al.*, 2008). MAPK activation results in phosphorylation of both nuclear transcription factors and other cytosolic targets that lead to changes in transcription, translation, cell motility, shape, proliferation, and differentiation (Derkinderen *et al.*, 2003; Lefkowitz and Shenoy, 2005). Furthermore, in response to prolonged activation, CB<sub>1</sub> receptor signalling is subject to regulation via receptor desensitization and internalization (Hsieh *et al.*, 1999; Jin *et al.*, 1999; Roche *et al.*, 1999; Marchese *et al.*, 2008). Desensitization is thought to result from phosphorylation of specific residues by GPCR kinases resulting in uncoupling of the receptor from G-protein signalling complexes. In contrast, internalization occurs via translocation of the receptor by endocytotic machinery to endosomes. Internalized receptors are subsequently recycled to the plasma membrane or degraded.

'Spice' is an herbal blend, marketed primarily in Europe for its cannabis-like effects and as an alternative to marijuana. A recent report used gas chromatography/mass spectrometry to analyse a number of different 'Spice' preparations as well as competing products from other manufacturers (Auwarter

*et al.*, 2009). Interestingly, these herbal 'Spice' blends contained diverse synthetic cannabinoid additives. Common among the different preparations was JWH018, a cannabinoid agonist from the aminoalkylindole family (Figure 1A). JWH018 has been shown to have a binding affinity for CB<sub>1</sub> receptors in the low nanomolar range (~9 nM) (Huffman *et al.*, 1994; Showalter *et al.*, 1996; Chin *et al.*, 1999; Aung *et al.*, 2000). In CB<sub>1</sub> receptor expressing CHO cells, JWH018 inhibits forskolin-stimulated cAMP production with an EC<sub>50</sub> of 14.7 nM with a maximal inhibition of 79% (Chin *et al.*, 1999). Beyond this there has been no report of the effect of JWH018 on CB<sub>1</sub> receptor-mediated cellular signalling. There has been a single report to date of the behavioural effects of JWH018 treatment. Wiley *et al.* (1998) found that JWH018 produced the tetrad of behaviours classically associated with cannabinoids (analgesia, catalepsy, hypomotility and hypothermia), having ED<sub>50</sub> values ranging from a low of 0.09 mg·kg<sup>-1</sup> for analgesia to a high of 1.47 mg·kg<sup>-1</sup> for hypothermia in the rodent model, suggesting that JWH018 activated CB<sub>1</sub> receptors *in vivo*.


Based on previous findings, we thought it was likely that JWH018 would act as an agonist in other CB<sub>1</sub> receptor-mediated signalling pathways and sought to characterize its ability to function as such. We examined the effect of JWH018 on neurotransmission and ERK1/2 MAPK activation and its ability to produce CB<sub>1</sub> receptor internalization. We found that JWH018 is both a potent and an efficacious CB<sub>1</sub> receptor agonist in each of these areas, actions that likely explain the ability of 'Spice' preparations to produce marijuana-like effects.

## Methods

### *Cell culture and transfection*

All animal care and experimental procedures used in this study were approved by the Animal Care Committees of Indiana University and conform to the Guidelines of the National Institutes of Health on the Care and Use of Animals. Mouse (CD1 strain and GAD67-GFP) hippocampal neurons isolated from the CA1–CA3 region were cultured on microislands as previously described (Furshpan *et al.*, 1976; Bekkers and Stevens, 1991). No significant differences were found between neurons isolated from either strain at any drug concentration tested, so the data from both strains were pooled. Neurons were obtained from animals (at postnatal day 0–2, killed via rapid decapitation without anaesthesia) and plated onto a feeder layer of hippocampal astrocytes that had been laid down previously (Levison and McCarthy, 1991). Cultures were grown in high-glucose (20 mM) minimum essential media containing 10% horse serum, without mitotic inhibitors and used for recordings after 8 days in culture and for no more than 3 h after removal from culture medium (Straiker and Mackie, 2005). All electrophysiological experiments were performed exclusively on excitatory neurons. All drugs were tested on cells from at least two different preparations.

Cell lines stably expressing CB<sub>1</sub> receptors were made as previously described (Brown *et al.*, 2002; Daigle *et al.*, 2008). Stable clones, uniformly expressing CB<sub>1</sub> receptors, were expanded and used for internalization and MAPK assays. Cells



**Figure 1** JWH018 decreases neurotransmitter release by activating presynaptic CB<sub>1</sub> receptors. (A) Chemical structure of JWH018. (B) JWH018 concentration-dependently decreased EPSC charge ( $n = 8$  to 14 for each concentration tested). This inhibitory effect was reversed by 1  $\mu$ M rimonabant ( $n = 12$ ) and was absent in CB<sub>1</sub><sup>-/-</sup> neurons ( $n = 5$ ). (C) A representative experiment showing a time course of EPSC inhibition by 1 nM JWH018 followed by 100 nM JWH018. 1  $\mu$ M rimonabant reversed JWH018 inhibition. Inset shows representative traces for three indicated time points. JWH018 (100 nM) significantly increases the paired-pulse ratio (right: representative traces of paired pulses, numbers indicate order of pulses) (D) and decreases miniature EPSC frequency (E) without affecting miniature EPSC amplitude (F), indicative of decreasing the probability of neurotransmitter release by acting at a presynaptic site ( $n = 5$  to 8 for each). Inset scale bars: 1 nA and 10 ms. (G) Coefficient of variation analysis demonstrating that  $r < \pi < 1$ , which is consistent with a presynaptic site for JWH018's action resulting in synaptic depression. Values are presented as mean  $\pm$  SEM where applicable. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.0001$ . CB<sub>1</sub>, cannabinoid receptor 1; EPSC, excitatory postsynaptic current.

were grown in Dulbecco's modified Eagle's media with 10% fetal bovine serum and penicillin/streptomycin (GIBCO, Carlsbad, CA, USA) at 37°C in 5% CO<sub>2</sub>.

#### Electrophysiology

When a single neuron is grown on a small island of permissive substrate, it forms synapses – or 'autapses' – onto itself.

All experiments were performed on isolated autaptic neurons. Whole-cell, voltage-clamp recordings from autaptic neurons were carried out at room temperature using an Axopatch 200A amplifier (Axon Instruments, Burlingame, CA, USA). The extracellular solution contained (mM) NaCl 119, KCl 5, CaCl<sub>2</sub> 2, MgCl<sub>2</sub> 1, glucose 30 and HEPES 20. Continuous flow of solution through the bath chamber (2 mL·min<sup>-1</sup>) ensured rapid drug application and clearance. Drugs were typically

prepared as a stock then diluted into extracellular solution at their final concentration and used on the same day. Recording pipettes of 1.8–4 M $\Omega$  were filled with solution containing (mM): potassium gluconate 121.5, KCl 17.5, NaCl 9, MgCl<sub>2</sub> 1, HEPES 10, EGTA 0.2, MgATP 2 and LiGTP 0.5. Access resistance was monitored and only cells with a stable access resistance were included for data analysis.

The membrane potential was held at  $-70$  mV and excitatory postsynaptic currents (EPSCs) were evoked every 20 s by triggering an unclamped action current with a 1.0 ms depolarizing step. The resultant evoked waveform consisted of a brief stimulus artefact (i.e. a large downward spike representing inward sodium currents) followed by the slower EPSC. The size of the recorded EPSCs was calculated by integrating the evoked current to yield charge (in pC). Calculating the charge in this manner yields an indirect measure of the amount of neurotransmitter released while minimizing the effects of cable distortion on currents generated far from the site of the recording electrode (the soma). Data were acquired at a sampling rate of 5 kHz.

Depolarization suppression of excitation (DSE) is a process whereby depolarization of a neuron results in production of endocannabinoids and activation of presynaptic CB<sub>1</sub> receptors with a subsequent transient decrease of glutamate release and EPSC amplitude. Cultured autaptic neurons are heterogeneous, with some expressing CB<sub>1</sub> receptors and others not. The presence of DSE (which requires CB<sub>1</sub> receptors) was used as a marker for neuronal cannabinoid sensitivity. DSE was induced as previously described (Straiker and Mackie, 2005). Recordings were primarily (see results) made from cells that exhibited DSE.

To determine the site of JWH018's action on neurotransmission, paired pulse ratio analysis, miniature EPSC (mEPSC) recordings and measurements of changes in the coefficient of variation (CV) were performed. Paired pulse ratios were calculated as the charge of the first of two pulses (60 ms interval) divided by the second. Ratios smaller than 1.0 were interpreted as paired-pulse depression. Analysis was performed only on neurons that displayed paired-pulse depression under basal conditions, indicating neurons with high probabilities of release. Ratios were calculated for each neuron under basal and drug-treated conditions.

mEPSC analysis was performed by measuring the frequency (Hz) and amplitude (pA) of mEPSC events under basal and drug-treated conditions. mEPSC events were analysed without knowledge of the treatments, and then the data from each condition were pooled. Plots of cumulative probability and graphs of mean mEPSC frequency and amplitude were made from this pooled data.

CV analysis was performed as described in Faber and Korn (1991) and Shen *et al.* (1996). Means and coefficients of variation were calculated from 6 to 20 sweeps from basal and drug-treated conditions.  $\pi$  and  $r$  were calculated for each individual experiment and means  $\pm$  SEM were calculated for each. A presynaptic site of drug action leading to synaptic depression was deduced if  $r < \pi < 1$ .

#### MAPK and receptor internalization assays

MAPK activation was analysed as previously described (Daigle *et al.*, 2008), with a few modifications. HEK293 cells stably

expressing CB<sub>1</sub> receptors were plated onto poly-D lysine-coated 96 well plates (Corning, Corning, NY) and allowed to grow to  $\sim 95\%$  confluency. The following day the growth media was removed and replaced with serum-free growth media and the cells were incubated overnight. The cells were washed once with HEPES buffered saline (HBS; 130 mM NaCl, 5.4 mM KCl, 1.8 mM MgCl<sub>2</sub>, and 10 mM HEPES, pH 7.5) containing 0.2 mg·mL<sup>-1</sup> bovine serum albumin (BSA). Drug containing solutions were made in the HBS/BSA solution and added to the wells at appropriate time points. Following drug incubation, the wells were emptied and ice-cold 4% paraformaldehyde was added immediately to each well, and the plates were placed on ice for 15 min followed by 30 min at room temperature. The paraformaldehyde was then removed and at least 100  $\mu$ L of ice-cold methanol was added to each well and the plate was incubated at  $-20^{\circ}\text{C}$  for at least 20 min. For methanol incubation times shorter than an hour, an additional washing step was performed using phosphate-buffered saline (PBS; 137 mM NaCl, 10 mM NaH<sub>2</sub>PO<sub>4</sub>, 2.7 mM KCl, pH 7.4) containing 0.1% Triton-X 100 for 25 min. The methanol or PBS/Triton-X 100 was replaced with a blocking solution of Tris-buffered saline (TBS; 137 mM NaCl, 10 mM Tris, pH 7.4) containing 5 mg·mL<sup>-1</sup> BSA and incubated for at least 1 h at room temperature. The blocking solution was then removed and replaced by a blocking solution containing rabbit anti-phospho-ERK1/2 MAPK antibody (1:200) (Cell Signaling Technologies Inc., Danvers, MA) and was allowed to shake overnight at 4 $^{\circ}\text{C}$  or for 3 h at room temperature. The antibody solution was removed and the plates were washed five times with TBS containing 0.05% Tween-20 (TBST) for 5–15 min each time. A blocking solution containing an IRDye conjugated anti-rabbit IgG antibody [either donkey anti-rabbit IR800 (1:500 dilution) or goat anti-rabbit IgG IR680 (1:200) antibody (LI-COR Biosciences, Lincoln, NE)] was added and allowed to shake for 1 h at room temperature. The plates were then washed five times with TBST, 5–15 min each time, followed with a final quick single rinse in distilled water. The plates were patted dry and then scanned using a LI-COR Odyssey. Integrated intensities were used for each well. The amount of MAPK activation was calculated as the average integrated intensities of the drug-treated wells divided by the average integrated intensities of the untreated wells and are expressed as percentages.

The extent of CB<sub>1</sub> receptor internalization was analysed as previously described in (Daigle *et al.*, 2008). The extent of internalization was calculated as the average integrated intensities of the drug-treated wells divided by the average integrated intensities of the untreated wells and are expressed as percentages.

#### Data analysis

Data are reported as mean  $\pm$  SEM (except EC<sub>50</sub>, IC<sub>50</sub> and t<sub>1/2</sub> data, which are reported as mean  $\pm$  95% CI). Non-linear regression was used to fit the concentration response curves and the time course of internalization. Paired Student's *t*-tests were used to evaluate the effect of drugs on paired pulse ratios and mEPSC data and unpaired Student's *t*-tests were used on all other comparison analyses. Statistical significance is indicated as follows: \*\*\**P* < 0.0001, \*\**P* < 0.01, and \**P* < 0.05. All

graphs and statistical analyses were generated using GraphPad Prism 4.0 software (Hearne Scientific Software, Chicago, IL, USA).

### Materials

Drugs and reagents were purchased from Tocris Cookson (Ellisville, MO, USA), Cayman Chemical (Ann Arbor, MI, USA) or Sigma-Aldrich (St Louis, MO, USA). JWH018 was synthesized as described by Huffman *et al.* (1994). Heterozygote (CB1+/-) mice to establish our colony were generously provided by Dr Catherine Ledent (University of Brussels, Belgium; Reibaud *et al.*, 1999). GAD67-GFP mice generated by Dr Yuchio Yanagawa (Gunman University, Gunma, Japan; Tamamaki *et al.*, 2003) were provided by Dr Albert Berger (University of Washington, Seattle, WA, USA) and used with Dr Yanagawa's permission. Rimonabant (SR141716) was obtained from the National Institute of Drug Abuse drug supply.

### Results

#### *JWH018 decreases the probability of neurotransmitter release via CB<sub>1</sub> receptor activation*

In the CNS, CB<sub>1</sub> receptors are predominately located on axon terminals. Upon agonist binding, CB<sub>1</sub> receptor activation leads to a reduced probability of neurotransmitter release. To see whether JWH018 acts similarly to other CB<sub>1</sub> receptor agonists in this regard, we recorded EPSCs from glutamatergic autaptic neuron cultures in the presence and absence of JWH018. Autaptic neuronal cultures are a well-characterized preparation that uses a single electrode to both stimulate and record EPSCs (Bekkers and Stevens, 1991). We found that JWH018 potently inhibited EPSCs (Figure 1B,C) in a concentration-dependent manner with a mean IC<sub>50</sub> of 14.9 nM (4.8–45.9 nM; 95% CI) and a maximal inhibition of 48.0 ± 4.2% of control at 1 μM. As seen in Figure 1C, the inhibition by JWH018 was poorly reversed upon washout of the drug. However, the effect of JWH018 (1 μM) on EPSC charge was fully reversed by 1 μM rimonabant (Figure 1C), a CB<sub>1</sub> antagonist, suggesting that the limited reversal is due to persistent receptor occupancy. The effect of JWH018 on EPSCs was examined in neurons that exhibited DSE, a transient reduction in EPSC size that results from depolarization. DSE is a well-described process that has been shown to be dependent on cannabinoid signalling molecules including CB<sub>1</sub> receptors. We found that in neurons that did not exhibit DSE (and are likely to lack CB<sub>1</sub> receptors); JWH018 had no effect on EPSC size (data not shown). To confirm that the effect observed with JWH018 was indeed due to action at CB<sub>1</sub> receptors, we treated neurons from CB<sub>1</sub> receptor null mice with 1 μM JWH018. In wild-type neurons, 1 μM JWH018 treatment reduced EPSC size (Figure 1B), whereas in the CB<sub>1</sub> receptor null neurons, 1 μM JWH018 had no effect (Figure 1B). To establish that JWH018 was operating through presynaptic CB<sub>1</sub> receptors to decrease the probability of neurotransmitter release, we performed paired-pulse analysis under baseline conditions and during JWH018 treatment. Figure 1D shows that 100 nM JWH018 treatment significantly increases the paired-pulse ratio ( $P = 0.0056$ ), suggesting a presynaptic site of


action. To further validate a presynaptic site of action we recorded mEPSCs before and after JWH018 treatment. Treatment of neurons with 100 nM JWH018 decreased mEPSC frequency (Figure 1E;  $P = 0.031$ ) but did not significantly alter mEPSC amplitude (Figure 1F;  $P = 0.19$ ). CV analysis further supported a presynaptic locus of JWH018's action (Figure 1G). 1 μM JWH018 treatment gave a mean value for  $r$  of  $0.26 \pm 0.049$  and a mean  $\pi$  value of  $0.54 \pm 0.41$ . All of these measures are consistent with a presynaptic site of drug action resulting in synaptic depression. In summary, JWH018 reduced the probability of glutamate release in autaptic neurons by acting at presynaptic CB<sub>1</sub> receptors in a concentration-dependent fashion.

#### *JWH018 activated MAPK*

In addition to modulating neurotransmission and cAMP levels, CB<sub>1</sub> receptor signalling activates MAPK activity in cultured cells (Bouaboula *et al.*, 1995; Daigle *et al.*, 2008) and neurons (Derkinderen *et al.*, 2003). This is conveniently detected by increased phosphorylation of ERK1/2. As JWH018 activated other CB<sub>1</sub> receptor-mediated signalling pathways we hypothesized that JWH018 would also activate ERK1/2. HEK293 cells stably expressing CB<sub>1</sub> receptors were treated with either JWH018 or WIN55,212, another well-characterized, efficacious CB<sub>1</sub> receptor agonist. The time course of ERK1/2 MAPK activation for 100 nM of each drug was determined. Maximal activation occurred between 5 and 10 min of treatment with both drugs (Figure 2A). Interestingly, at both 5 and 7.5 min JWH018 was a more efficacious activator of ERK1/2 phosphorylation. A concentration-response analysis was performed at the 7.5-min time point (Figure 2B). JWH018 was more potent with an EC<sub>50</sub> of 4.4 nM (1.6–12.5 nM) compared with WIN55,212, which had an EC<sub>50</sub> of 69.9 nM (37.2–131.4 nM). The two agonists had similar maximal effects at 1 μM, although JWH018 was more potent than WIN55,212. In addition, to ensure that the MAPK activation observed from these drug treatments required CB<sub>1</sub> receptors, the cells were treated with rimonabant and then agonist. Here, 1 μM rimonabant prevented the activation of MAPK by either 1 μM JWH018 or WIN 55,212 (JWH018: 97.1 ± 5.7%, WIN55,212: 86.8 ± 5.8%, NS vs. untreated). Thus, in addition to its inhibition of adenylyl cyclase and neurotransmission, JWH018 also serves as a potent agonist of CB<sub>1</sub> receptor-mediated ERK1/2 MAPK activation.


#### *JWH018-induced robust CB<sub>1</sub> receptor internalization*

Following prolonged exposure to agonists, many GPCRs undergo internalization. This is thought to be a means whereby the cell can control its spatial and temporal response to receptor agonists (Drake *et al.*, 2006; Marchese *et al.*, 2008). CB<sub>1</sub> receptor internalization has been described in response to numerous cannabinoid drugs (Hsieh *et al.*, 1999; Coutts *et al.*, 2001; Daigle *et al.*, 2008). We hypothesized that based on previous experiments, if JWH018 was acting as a CB<sub>1</sub> receptor agonist, long-term treatment of CB<sub>1</sub>



**Figure 2** JWH018 activation of CB<sub>1</sub> receptors stimulated ERK1/2 MAPK phosphorylation. (A) 100 nM JWH018 treatment of CB<sub>1</sub> expressing HEK293 cells, transiently increased ERK1/2 phosphorylation, similar to the time course of 100 nM WIN55,212 activation (4–6 replicate samples from 7–15 independent experiments). (B) CB<sub>1</sub> receptor expressing HEK293 cells were treated for 7.5 min with increasing concentrations of JWH018 and WIN55,212. JWH018 activated ERK1/2 in a concentration-dependent manner and was more potent than WIN55,212 (4–6 replicate samples from 8–15 independent experiments). 1 μM rimonabant reversed the effect of JWH018 and WIN55,212 on ERK1/2 activation. Values are presented as mean ± SEM. \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.0001: significantly different from WIN55,212 treatment. ####*P* < 0.0001: significantly different from rimonabant treatment. CB<sub>1</sub>, cannabinoid receptor 1; MAPK, mitogen activated protein kinase.

receptor expressing cells with JWH018 would lead to profound receptor internalization. Figure 3A shows the time course of CB<sub>1</sub> receptor internalization in HEK293 cells stably expressing the receptor. After 3 h of exposure to 100 nM JWH018, 38.1 ± 3.1% of the CB<sub>1</sub> receptors present at the plasma membrane under basal conditions remained on the cell surface (61.9% internalization). We again used an equal concentration of WIN55,212 as a control, in which 3 h of treatment led to 58.7 ± 3.2% internalization (NS vs. JWH018). JWH018 induced a much more rapid internalization than WIN55,212 with a *t*<sub>1/2</sub> of 17.3 min (13.6–23.7 min) compared with 39.6 min (26.2–80.9 min) for WIN 55,212. Each drug caused similar maximal levels of internalization reaching plateaus of approximately 60% internalization (JWH018 = 40.7 ± 1.8% of basal levels; WIN55,212 = 41.1 ± 4.9% of basal levels). We performed concentration-response analysis of internalization using a 2 h treatment period (Figure 3B). JWH018 and WIN55,212 had similar maximal effects reaching 54.7 ± 3.4% and 56.0 ± 3.0% internaliza-



**Figure 3** JWH018 induced CB<sub>1</sub> receptor internalization. (A) In CB<sub>1</sub> receptor expressing HEK293 cells, 100 nM JWH018 treatment resulted in robust receptor internalization that was quicker than that induced by 100 nM WIN55,212 (4–6 replicate samples from 5–10 independent experiments). (B) Following 2 h of exposure to increasing concentrations of each drug, CB<sub>1</sub> receptors were internalized in a concentration-dependent manner with JWH018 being more potent than WIN55,212 (4–6 replicate samples from 5–10 independent experiments). 1 μM rimonabant, a CB<sub>1</sub> receptor antagonist, prevented receptor internalization due to JWH018 or WIN55,212 treatment. Values are presented as mean ± SEM. \**P* < 0.05, \*\**P* < 0.01: significantly different from WIN55,212 treatment. ####*P* < 0.0001: significantly different from rimonabant treatment. CB<sub>1</sub>, cannabinoid receptor 1.

tion at 1 μM concentrations respectively (NS). However, JWH018 was once again the more potent with an EC<sub>50</sub> of 2.8 nM (1.2–6.3 nM) compared with WIN55,212, which had an EC<sub>50</sub> of 19.4 nM (8.5–44.4 nM). 1 μM rimonabant prevented internalization by 1 μM of either drug (JWH018: 91.5 ± 3.8%, WIN55,212: 103.3 ± 3.3%, NS vs. untreated). Therefore, prolonged exposure to JWH018 leads to robust CB<sub>1</sub> receptor internalization, as did treatment with WIN55,212, although JWH018 was more potent and caused faster internalization.

## Discussion

JWH018 is a common synthetic additive found in diverse preparations of the herbal blend known as 'Spice' (Auwarter *et al.*, 2009). JWH018 was first synthesized during an analysis of cannabimimetic indole structures that aimed to design new indoles with effects comparable with those of natural cannabinoids such as THC (Huffman *et al.*, 1994). Relatively little characterization of this ligand has been performed. It has been reported to have a high affinity for CB<sub>1</sub> receptor with a

$K_i$  of approximately 9 nM (Huffman *et al.*, 1994; Showalter *et al.*, 1996; Chin *et al.*, 1999; Aung *et al.*, 2000) and to inhibit adenylyl cyclase with an  $IC_{50}$  of 14.7 nM and a 79% maximal inhibition. Beyond these preliminary studies, there have been no other investigations of JWH018's effect on CB<sub>1</sub> receptor-mediated signalling. In light of its presence in 'Spice' and its cannabinoid-like psychoactivity, we have examined the effects of this compound on neurotransmission, MAPK activity, and CB<sub>1</sub> receptor internalization.

CB<sub>1</sub> receptor activation may suppress neurotransmission and neuronal excitability (Kano *et al.*, 2009). Here we have used autaptic excitatory hippocampal neurons as a well-characterized model system of cannabinoid-mediated effects on neurotransmission (Straiker and Mackie, 2005; 2007). We found that JWH018 potently inhibits glutamate release in these neurons in a concentration-dependent fashion. The effect of JWH018 on synaptic transmission was due to its action at CB<sub>1</sub> receptors as JWH018 had no effect on EPSC in neurons cultured from CB<sub>1</sub> receptor knockout mice, and rimonabant, a CB<sub>1</sub> receptor antagonist, blocked its effect in wild-type neurons. Furthermore, JWH018 is very likely acting at presynaptic CB<sub>1</sub> receptors based on its ability to increase the paired-pulse ratio, to decrease the frequency of mEPSCs without affecting mEPSC amplitude and to increase the CV. Cannabinoid agonists sometimes do not decrease mEPSC frequency at glutamatergic synapses (Yamasaki *et al.*, 2006). However, our results here are consistent with most reports that find cannabinoid suppression of mEPSC frequency (see Misner and Sullivan, 1999; Sullivan, 1999; Morisset and Urban, 2001; Robbe *et al.*, 2001; Derbenev *et al.*, 2004), and the difference may lie in the brain region, cell type or culture conditions used. The effect of JWH018 on neurotransmission is both potent and effective with an  $IC_{50}$  of 14.9 nM and a maximal inhibition to 48.0% of control at 1  $\mu$ M. JWH018 exhibited effects comparable with those of WIN55,212 (Straiker and Mackie, 2005). In light of the internalization data, JWH018's effect on neurotransmission may potentially be influenced by receptor desensitization or internalization. We cannot make any certain conclusions based on our data here as to whether desensitization had an effect on our recordings. Internalization is unlikely to play a role as little CB<sub>1</sub> receptor internalization would occur at room temperature during the 20–30 min of recording. In summary, JWH018 inhibits synaptic transmission as a potent and efficacious CB<sub>1</sub> receptor agonist.

Based on the earlier studies discussed above and our finding that JWH018 inhibited synaptic neurotransmission, it seemed probable that JWH018 would exhibit signalling effects similar to those of other CB<sub>1</sub> receptor agonists. However, it was important to test this hypothesis as JWH018 may exhibit a different functional selectivity relative to other cannabinoid agonists (Urban *et al.*, 2007). ERK1/2 MAPK activation is a typical consequence of CB<sub>1</sub> receptor stimulation (Bouaboula *et al.*, 1995; Daigle *et al.*, 2008). We have demonstrated that as observed with other cannabinoid ligands such as WIN55,212, JWH018 also stimulated ERK1/2 MAPK activation in a concentration-dependent manner. This ERK1/2 activation had the typically observed rapid time course (Daigle *et al.*, 2008) reaching a peak level of activation between 5 and 10 min. Comparing JWH018 with WIN55,212, we found that

despite having a similar time course of activation, JWH018 was more potent with an  $EC_{50}$  of 4.4 nM compared with WIN55,212 with an  $EC_{50}$  of 69.9 nM. Both were similarly efficacious.

CB<sub>1</sub> receptor internalization has been reported to occur in response to a number of different cannabinoid ligands in a number of cell types (Hsieh *et al.*, 1999; Coutts *et al.*, 2001; Daigle *et al.*, 2008). Here we found that JWH018, consistent with its ability to act as a CB<sub>1</sub> receptor agonist, produces robust CB<sub>1</sub> receptor internalization that is rapid ( $t_{1/2}$  = 17.3 min), potent ( $EC_{50}$  = 2.8 nM) and efficacious (38.1% of basal surface levels at 3 h with 100 nM). We found it more potent and to induce internalization more rapidly than WIN55,212 ( $EC_{50}$  = 19.4 nM and  $t_{1/2}$  = 39.6 min). However, the two drugs are of similar efficacy (JWH018 – plateaus at 40.7% and WIN55,212 – plateaus at 41.1% of basal surface levels). Thus, prolonged engagement of CB<sub>1</sub> receptors by JWH018, as with many other CB<sub>1</sub> receptor agonists, leads to profound cellular adaptations that may serve to decrease cellular sensitivity to the drug. Interpretation of the effects of chronic JWH018 on behaviour must consider its ability to produce cellular desensitization and tolerance.

Previously we found that in autaptic neuronal cultures, THC did not inhibit EPSCs but rather, it antagonized inhibition by both WIN55,212 and 2-arachidonyl glycerol (2-AG) (Straiker and Mackie, 2005). Similar results have been reported in non-autaptic hippocampal cultures (Roloff and Thayer, 2009). In these systems, THC effectively acts as an antagonist with short-term treatment but desensitizes CB<sub>1</sub> receptor signalling with long-term treatment. Since THC, which is the principal psychoactive component of marijuana, is a low-efficacy CB<sub>1</sub> receptor agonist, we speculated that it prevents full CB<sub>1</sub> receptor activation by the endocannabinoid 2-AG and mimics the effects of the low-efficacy endocannabinoid anandamide (Straiker and Mackie, 2005). This was a provocative hypothesis to explain the psychoactive effects of marijuana. However, in light of the findings from this study, this hypothesis must be revised. 'Spice' apparently produces marijuana-like psychoactivity when smoked (Auwarter *et al.*, 2009). However, it has not been reported to contain THC but rather contains at least one potent and efficacious CB<sub>1</sub> receptor agonist. Therefore, if the cognitive effects of 'Spice' are due to JWH018, our proposition that the psychoactivity of THC may in part be due to the antagonism of 2-AG activation of CB<sub>1</sub> receptors requires rethinking. The effects we previously observed with THC may be unique to the cultured neuron preparation itself. In cultures, the number of CB<sub>1</sub> receptors or their coupling may be limited, which will cause a low-efficacy agonist to act as an antagonist. In contrast, these factors may not be limiting for CB<sub>1</sub> receptors expressed in brain, and a low-efficacy agonist such as THC may act as an agonist *in vivo*. Alternatively, as was suggested in a recent study, the difference may also lie in the firing rate of the neurons, which can influence a neuron's response to THC (Roloff and Thayer, 2009). Furthermore, it may be a possibility that there are other uncharacterized additives in 'Spice' that influence neurotransmission. Despite these uncertainties, it is clear that JWH018 potently inhibits neurotransmission, with an efficacy comparable with other synthetic cannabinoids such as WIN55,212.

Among the different preparations of 'Spice' that were analysed by Auwarter *et al.* (2009), JWH018 was a frequent additive. Here we have demonstrated that JWH018 treatment has cellular effects similar to those of other efficacious cannabinoid agonists such as WIN55,212. We have found that JWH018 is a more potent CB<sub>1</sub> receptor agonist than WIN55,212, although of similar efficacy. This is consistent with the reports that 'Spice' has marijuana-like effects when smoked. While Auwarter *et al.* found that JWH018 was not the most abundant of the additives present in various spice preparations, its high potency suggests that it will produce behavioural effects in humans. The selectivity of JWH018 for CB<sub>1</sub> receptors is low: JWH018 has a  $K_i$  of about 9 nM at CB<sub>1</sub> receptors and a  $K_i$  of about 3 nM at CB<sub>2</sub> receptors (Huffman *et al.*, 1994; Chin *et al.*, 1999; Aung *et al.*, 2000). While the effects we observed are clearly due to CB<sub>1</sub> receptor activation, the potential role of CB<sub>2</sub> receptors in the effects of 'Spice' requires further study. This study has focused on JWH018; however, different preparations of 'Spice' apparently contain diverse synthetic additives such as a modified version of CP47,497 (extending the dimethylheptyl side chain to dimethyloctyl), a cannabinoid ligand, that may also act as agonists at CB<sub>1</sub> receptors but so far remain uncharacterized and HU210 (Huffman *et al.*, 2008; Auwarter *et al.*, 2009). It is likely that these additional compounds might also contribute to the behavioural and subjective effects produced by smoking 'Spice', and their different pharmacologies might cause different preparations of 'Spice' to vary in their psychoactivity or health effects. Investigation into these additional synthetic additives requires further attention. Despite these caveats, we have shown that JWH018 has profound CB<sub>1</sub> receptor-mediated effects on cellular signalling and neurotransmission, which are likely to have a significant impact on cognitive function. Thus, 'Spice', which is marketed as a 'natural' herbal blend, is actually a vehicle of delivery for at least one very potent synthetic CB<sub>1</sub> receptor agonist, and its presence is likely to account for the psychoactive effects produced when 'Spice' is smoked.

## Acknowledgements

Supported by DA11322, DA21696, DA024122, DA009158 and DA003590.

## Conflict of interest

None.

## References

- Alexander SP, Mathie A, Peters JA (2008). Guide to receptors and channels (GRAC), 3rd edition. *Br J Pharmacol* **153** (Suppl. 2): S1–S209.
- Ameri A, Wilhelm A, Simmet T (1999). Effects of the endogenous cannabinoid, anandamide, on neuronal activity in rat hippocampal slices. *Br J Pharmacol* **126**: 1831–1839.
- Aung MM, Griffin G, Huffman JW, Wu M, Keel C, Yang B *et al.* (2000). Influence of the N-1 alkyl chain length of cannabimimetic indoles upon CB(1) and CB(2) receptor binding. *Drug Alcohol Depend* **60**: 133–140.
- Auwarter V, Dresen S, Weinmann W, Muller M, Putz M, Ferreiros N (2009). 'Spice' and other herbal blends: harmless incense or cannabinoid designer drugs? *J Mass Spectrom* **44**: 832–837.
- Bekkers JM, Stevens CF (1991). Excitatory and inhibitory autaptic currents in isolated hippocampal neurons maintained in cell culture. *Proc Natl Acad Sci USA* **88**: 7834–7838.
- Bouaboula M, Poinot-Chazel C, Bourrie B, Canat X, Calandra B, Rinaldi-Carmona M *et al.* (1995). Activation of mitogen-activated protein kinases by stimulation of the central cannabinoid receptor CB1. *Biochem J* **312** (Pt 2): 637–641.
- Brown SM, Wager-Miller J, Mackie K (2002). Cloning and molecular characterization of the rat CB2 cannabinoid receptor. *Biochim Biophys Acta* **1576**: 255–264.
- Chevalere V, Takahashi KA, Castillo PE (2006). Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu Rev Neurosci* **29**: 37–76.
- Chin CN, Murphy JW, Huffman JW, Kendall DA (1999). The third transmembrane helix of the cannabinoid receptor plays a role in the selectivity of aminoalkylindoles for CB2, peripheral cannabinoid receptor. *J Pharmacol Exp Ther* **291**: 837–844.
- Costa B (2007). On the pharmacological properties of delta9-tetrahydrocannabinol (THC). *Chem Biodivers* **4**: 1664–1677.
- Coutts AA, Anavi-Goffer S, Ross RA, MacEwan DJ, Mackie K, Pertwee RG *et al.* (2001). Agonist-induced internalization and trafficking of cannabinoid CB1 receptors in hippocampal neurons. *J Neurosci* **21**: 2425–2433.
- Daigle TL, Kearn CS, Mackie K (2008). Rapid CB1 cannabinoid receptor desensitization defines the time course of ERK1/2 MAP kinase signaling. *Neuropharmacology* **54**: 36–44.
- Derbenev AV, Stuart TC, Smith BN (2004). Cannabinoids suppress synaptic input to neurons of the rat dorsal motor nucleus of the vagus nerve. *J Physiol* **559**: 923–938.
- Derkinderen P, Valjent E, Toutant M, Corvol JC, Enslen H, Ledent C *et al.* (2003). Regulation of extracellular signal-regulated kinase by cannabinoids in hippocampus. *J Neurosci* **23**: 2371–2382.
- Drake MT, Shenoy SK, Lefkowitz RJ (2006). Trafficking of G protein-coupled receptors. *Circ Res* **99**: 570–582.
- Faber DS, Korn H (1991). Applicability of the coefficient of variation method for analyzing synaptic plasticity. *Biophys J* **60**: 1288–1294.
- Furshpan EJ, MacLeish PR, O'Lague PH, Potter DD (1976). Chemical transmission between rat sympathetic neurons and cardiac myocytes developing in microcultures: evidence for cholinergic, adrenergic, and dual-function neurons. *Proc Natl Acad Sci USA* **73**: 4225–4229.
- Gulyas AI, Cravatt BF, Bracey MH, Dinh TP, Piomelli D, Boscia F *et al.* (2004). Segregation of two endocannabinoid-hydrolyzing enzymes into pre- and postsynaptic compartments in the rat hippocampus, cerebellum and amygdala. *Eur J Neurosci* **20**: 441–458.
- Howlett AC (2002). The cannabinoid receptors. *Prostaglandins Other Lipid Mediat* **68–69**: 619–631.
- Howlett AC, Barth F, Bonner TI, Cabral G, Casellas P, Devane WA *et al.* (2002). International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol Rev* **54**: 161–202.
- Hsieh C, Brown S, Derleth C, Mackie K (1999). Internalization and recycling of the CB1 cannabinoid receptor. *J Neurochem* **73**: 493–501.
- Huestis MA, Gorelick DA, Heishman SJ, Preston KL, Nelson RA, Moolchan ET *et al.* (2001). Blockade of effects of smoked marijuana by the CB1-selective cannabinoid receptor antagonist SR141716. *Arch Gen Psychiatry* **58**: 322–328.
- Huffman JW, Dong D, Martin BR, Compton DR (1994). Design, synthesis and pharmacology of cannabimimetic indoles. *Bioorg Med Chem Lett* **4**: 563–566.



- Huffman JW, Thompson AL, Wiley JL, Martin BR (2008). Synthesis and pharmacology of 1-deoxy analogs of CP-47,497 and CP-55,940. *Bioorg Med Chem* **16**: 322–335.
- Jin W, Brown S, Roche JP, Hsieh C, Celver JP, Kovoor A *et al.* (1999). Distinct domains of the CB1 cannabinoid receptor mediate desensitization and internalization. *J Neurosci* **19**: 3773–3780.
- Kano M, Ohno-Shosaku T, Hashimoto Y, Uchigashima M, Watanabe M (2009). Endocannabinoid-mediated control of synaptic transmission. *Physiol Rev* **89**: 309–380.
- Kreitzer AC, Regehr WG (2001). Retrograde inhibition of presynaptic calcium influx by endogenous cannabinoids at excitatory synapses onto Purkinje cells. *Neuron* **29**: 717–727.
- Lefkowitz RJ, Shenoy SK (2005). Transduction of receptor signals by beta-arrestins. *Science* **308**: 512–517.
- Levison SW, McCarthy KD (1991). Characterization and partial purification of AIM: a plasma protein that induces rat cerebral type 2 astroglia from bipotential glial progenitors. *J Neurochem* **57**: 782–794.
- Mackie K (2005). Distribution of cannabinoid receptors in the central and peripheral nervous system. *Handb Exp Pharmacol* **168**: 299–325.
- Mackie K, Lai Y, Westenbroek R, Mitchell R (1995). Cannabinoids activate an inwardly rectifying potassium conductance and inhibit Q-type calcium currents in AtT20 cells transfected with rat brain cannabinoid receptor. *J Neurosci* **15**: 6552–6561.
- Marchese A, Paing MM, Temple BR, Trejo J (2008). G protein-coupled receptor sorting to endosomes and lysosomes. *Annu Rev Pharmacol Toxicol* **48**: 601–629.
- Matyas F, Watanabe M, Mackie K, Katona I, Freund TF (2007). Molecular architecture of the cannabinoid signaling system in the core of the nucleus accumbens. *Ideggyogy Sz* **60**: 187–191.
- Misner DL, Sullivan JM (1999). Mechanism of cannabinoid effects on long-term potentiation and depression in hippocampal CA1 neurons. *J Neurosci* **19**: 6795–6805.
- Monory K, Blaudzun H, Massa F, Kaiser N, Lemberger T, Schutz G *et al.* (2007). Genetic dissection of behavioural and autonomic effects of delta(9)-tetrahydrocannabinol in mice. *PLoS Biol* **5**: e269.
- Morisset V, Urban L (2001). Cannabinoid-induced presynaptic inhibition of glutamatergic EPSCs in substantia gelatinosa neurons of the rat spinal cord. *J Neurophysiol* **86**: 40–48.
- Nyiri G, Cserep C, Szabadits E, Mackie K, Freund TF (2005). CB1 cannabinoid receptors are enriched in the perisynaptic annulus and on preterminal segments of hippocampal GABAergic axons. *Neuroscience* **136**: 811–822.
- Ohno-Shosaku T, Maejima T, Kano M (2001). Endogenous cannabinoids mediate retrograde signals from depolarized postsynaptic neurons to presynaptic terminals. *Neuron* **29**: 729–738.
- Pertwee RG (2008). The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol* **153**: 199–215.
- Reibaud M, Obinu MC, Ledent C, Parmentier M, Bohme GA, Imperato A (1999). Enhancement of memory in cannabinoid CB1 receptor knock-out mice. *Eur J Pharmacol* **379**: R1–R2.
- Robbe D, Alonso G, Duchamp F, Bockaert J, Manzoni OJ (2001). Localization and mechanisms of action of cannabinoid receptors at the glutamatergic synapses of the mouse nucleus accumbens. *J Neurosci* **21**: 109–116.
- Roche JP, Bounds S, Brown S, Mackie K (1999). A mutation in the second transmembrane region of the CB1 receptor selectively disrupts G protein signaling and prevents receptor internalization. *Mol Pharmacol* **56**: 611–618.
- Roloff AM, Thayer SA (2009). Modulation of excitatory synaptic transmission by delta 9-tetrahydrocannabinol switches from agonist to antagonist depending on firing rate. *Mol Pharmacol* **75**: 892–900.
- Shen M, Piser TM, Seybold VS, Thayer SA (1996). Cannabinoid receptor agonists inhibit glutamatergic synaptic transmission in rat hippocampal cultures. *J Neurosci* **16**: 4322–4334.
- Showalter VM, Compton DR, Martin BR, Abood ME (1996). Evaluation of binding in a transfected cell line expressing a peripheral cannabinoid receptor (CB2): identification of cannabinoid receptor subtype selective ligands. *J Pharmacol Exp Ther* **278**: 989–999.
- Straiker A, Mackie K (2005). Depolarization-induced suppression of excitation in murine autaptic hippocampal neurones. *J Physiol* **569**: 501–517.
- Straiker A, Mackie K (2006). Cannabinoids, electrophysiology, and retrograde messengers: challenges for the next 5 years. *AAPS J* **8**: E272–E276.
- Straiker A, Mackie K (2007). Metabotropic suppression of excitation in murine autaptic hippocampal neurons. *J Physiol* **578** (Pt 3): 773–785.
- Sullivan JM (1999). Mechanisms of cannabinoid-receptor-mediated inhibition of synaptic transmission in cultured hippocampal pyramidal neurons. *J Neurophysiol* **82**: 1286–1294.
- Tamamaki N, Yanagawa Y, Tamioka R, Miyazaki J, Obata K, Kaneko T (2003). Green fluorescent protein expression and colocalization with calretinin, parvalbumin, and somatostatin in the GAD67-GFP knock-in mouse. *J Comp Neurol* **467**: 60–79.
- Taura F, Sirikantaramas S, Shoyama Y, Morimoto S (2007). Phytocannabinoids in Cannabis sativa: recent studies on biosynthetic enzymes. *Chem Biodivers* **4**: 1649–1663.
- Twitchell W, Brown S, Mackie K (1997). Cannabinoids inhibit N- and P/Q-type calcium channels in cultured rat hippocampal neurons. *J Neurophysiol* **78**: 43–50.
- Urban JD, Clarke WP, von Zastrow M, Nichols DE, Kobilka B, Weinstein H *et al.* (2007). Functional selectivity and classical concepts of quantitative pharmacology. *J Pharmacol Exp Ther* **320**: 1–13.
- Wiley JL, Compton DR, Dai D, Lainton JA, Phillips M, Huffman JW *et al.* (1998). Structure–activity relationships of indole- and pyrrole-derived cannabinoids. *J Pharmacol Exp Ther* **285**: 995–1004.
- Wilson RI, Nicoll RA (2001). Endogenous cannabinoids mediate retrograde signalling at hippocampal synapses. *Nature* **410**: 588–592.
- Yamasaki M, Hashimoto K, Kano M (2006). Miniature synaptic events elicited by presynaptic Ca<sup>2+</sup> rise are selectively suppressed by cannabinoid receptor activation in cerebellar Purkinje cells. *J Neurosci* **26**: 86–95.
- Yoshida T, Fukaya M, Uchigashima M, Miura E, Kamiya H, Kano M *et al.* (2006). Localization of diacylglycerol lipase-alpha around postsynaptic spine suggests close proximity between production site of an endocannabinoid, 2-arachidonoyl-glycerol, and presynaptic cannabinoid CB1 receptor. *J Neurosci* **26**: 4740–4751.